Author Index

Aderjan, R.
—— and Bogusz, M.
Nitroalkanes as a multidetector retention index scale for drug identification in gas chromatography 345
Andersson, K., see Levin, J.-O. 121
Angers, D. A.
——, Nadeau, P. and Mehuys, G. R.
Determination of carbohydrate composition of soil hydrolysates by high-performance liquid chromatography 444
Banks, M. A., see Porter, D. W. 311
Bartley, T. D., see Klein, M. L. 205
Battista, M.
——, Di Corcia, A. and Marchetti, M.
High-performance liquid chromatographic method for determining triazine herbicide residues in soil 233
Beger, J., see Voelkel, A. 51
Bergers, P. J. M., see Niessen, W. M. A. 243
Bidlingmeyer, B., see Chou, T.-Y. 169
Biernat, J. F., see Kostrowicki, J. 340
Blaas, W., see Tiebach, R. 372
Blessington, B.
—— and Crabb, N.
Chiral high-performance liquid chromatographic analysis of phenoxy herbicide mixtures 450
Bogusz, M., see Aderjan, R. 345
Bufalo, G., see Grasso, G. 411
Campbell, L. D., see Slominski, B. A. 285
Castello, G.
——, Timossi, A. and Gerbino, T. C.
Gas chromatographic separation of halogenated compounds on non-polar and polar wide bore capillary columns 129
Castranova, V., see Porter, D. W. 311
Čelap, M. B., see Vučković, G. 362
Čellár, P., see Matisová, E. 65
Chou, F. E., see Ito, Y. 382
Chou, T.-Y.
——, Gao, C.-X., Colgan, S. T., Krull, I. S., Dorschel, C. and Bidlingmeyer, B.
Polymeric benzotriazole reagent for the off-line high-performance liquid chromatographic derivatization of polyamines and related nucleophiles in biological fluids 169
Christie, W. W.
Separation of molecular species of triacylglycerols by high-performance liquid chromatography with a silver ion column 273
Colgan, S. T., see Chou, T.-Y. 169
Colote, S., see Liautard, J. 195
Corcia, A. Di, see Battista, M. 233
Crabb, N., see Blessington, B. 450
Cramer, S. M., see Phillips, M. W. 1
Cygan, A., see Kostrowicki, J. 340
Dalene, M., see Skarping, G. 293
Dentshev, T., see Stoev, G. 367
Di Corcia, A., see Battista, M. 233
Dimitrov, Chr., see Gavrilova, T. B. 73
Dorschel, C., see Chou, T.-Y. 169
Eganhouse, R. P., see Sherblom, P. M. 37
Erni, F., see Steuer, W. 253
Euerby, M. R.
Effect of differing thiols on the reversed-phase high-performance liquid chromatographic behaviour of *o*-phthaldialdehyde-thiol–amino acids 398
Fábián-Varga, K., see Gazdag, M. 95
Fang, Q. C., see Zhang, T.-Y. 185
Ferraz, C., see Liautard, J. 195
Fujita, M., see Saito, K. 387
Gao, C.-X., see Chou, T.-Y. 169
Gasparič, J.
Dual retention mechanism on commercial acetylcellulose as a stationary phase in planar chromatography 352
Gavrilova, T. B.
——, Vlasenko, E. V., Petsev, N., Topalova, I., Dimitrov, Chr. and Ivanov, S.
Gas chromatographic retention on carbon adsorbents coated with monolayers of polynuclear aromatic hydrocarbons 73
Gazdag, M.
——, Szepesi, G. and Fábián-Varga, K.
Selection of high-performance liquid chromatographic methods in pharmaceutical analysis. II. Optimization for selectivity in normal-phase systems 95
——, Szepesi, G. and Szeleczki, E.
Selection of high-performance liquid chromatographic methods in pharmaceutical analysis. I. Optimization for selectivity in reversed-phase chromatography 83
Gerbino, T. C., see Castello, G. 129
Grasso, G.
—— and Bufalo, G.
Application of ion chromatography to the analysis of chromium hydroxy salts 411
Greef, J. van der, see Niessen, W. M. A. 243
Hartwick, R. A., see Raglione, T. V. 157
Haynes, S., see Longmuir, K. J. 438
Horbaczewski, A., see Kostrowicki, J. 340

Horie, M., see Saito, K. 387
Horiuchi, K., see Suemitsu, R. 406
Hoshino, Y., see Saito, K. 387
Hulst, A. G., see Wils, E. R. J. 261
Husain, S.
——, Sastry, G. S. R., Prasada Raju, N. and Narasimha, R.
High-performance size-exclusion chromatography of oils and fats 317
Ibrahim, E. A.
—— and Suffet, I. H.
Freon FC-113, an alternative to methylene chloride for liquid–liquid extraction of trace organics from chlorinated drinking water 217
Ioffe, B. V., see Marinichev, A. N. 327
Ishii, J., see Terabayashi, T. 432
Ito, Y.
—— and Chou, F. E.
New high-speed counter-current chromatograph equipped with a pair of separation columns connected in series 382
——, see Zhang, T.-Y. 185
Ivanov, S., see Gavrilova, T. B. 73
Kallio, H.
—— and Shibamoto, T.
Direct capillary trapping and gas chromatographic analysis of bromomethane and other highly volatile air pollutants 392
Kaneshima, H., see Kanetoshi, A. 145
Kanetoshi, A.
——, Ogawa, H., Katsura, E., Kaneshima, H. and Miura, T.
Formation of polychlorinated dibenzo-*p*-dioxin from 2,4,4′-trichloro-2′-hydroxydiphenyl ether (Irgasan® DP300) and its chlorinated derivatives by exposure to sunlight 145
Karlsson, R.-M., see Levin, J.-O. 121
Katsura, E., see Kanetoshi, A. 145
Kaufman, A. E.
—— and Polymeropoulos, C. E.
Study of the injection process in a gas chromatograph split injection port 23
Kawanishi, Y., see Terabayashi, T. 432
Kennerly, D. A.
Two-dimensional thin-layer chromatographic separation of phospholipid molecular species using plates with both reversed-phase and argentation zones 425
Kimura, Y., see Tokuda, H. 109
Klein, M. L.
——, Bartley, T. D., Lai, P.-H. and Lu, H. S.
Structural characterization of recombinant consensus interferon-α 205
Koizumi, K.
——, Kubota, Y., Tanimoto, T. and Okada, Y.
Determination of cyclic glucans by anion-exchange chromatography with pulsed amperometric detection 303
Kostrowicki, J.
——, Luboch, E., Makuch, B., Cygan, A., Horbaczewski, A. and Biernat, J. F.
Macrocyclic polyfunctional Lewis bases. XII. Influence of complex formation on chromatographic migration 340
Krull, I. S., see Chou, T.-Y. 169
Krupčík, J., see Matisová, E. 65
Kubota, Y., see Koizumi, K. 303
Lai, P.-H., see Klein, M. L. 205
Leclercq, P. A., see Matisová, E. 65
Lee, Y.-W., see Zhang, T.-Y. 185
Levin, J.-O.
——, Andersson, K. and Karlsson, R.-M.
Solid sorbent sampling and chromatographic determination of glycidyl ethers in air 121
Liautard, J.
——, Colote, S., Ferraz, C., Sri-Widada, J., Pichot, A. and Liautard, J. P.
Analysis of double-stranded Poly(A) · Poly(U) molecules by reversed-phase high-performance liquid chromatography 195
Liautard, J. P., see Liautard, J. 195
Longmuir, K. J.
—— and Haynes, S.
Purification of long-chain, saturated, free fatty acids 438
Lu, H. S., see Klein, M. L. 205
Luboch, E., see Kostrowicki, J. 340
Makuch, B., see Kostrowicki, J. 340
Malinar, M. J., see Vučković, G. 362
Marchetti, M., see Battista, M. 233
Marinichev, A. N.
—— and Ioffe, B. V.
Theoretical aspects of the application of headspace analysis to the investigation of reaction kinetics in solutions 327
Martin, W. G., see Porter, D. W. 311
Matisová, E.
——, Moravcová, A., Krupčík, J., Čellár, P. and Leclercq, P. A.
Problems with the reproducibility of retention data on capillary columns with hydrocarbon C_{87} as the stationary phase 65
Mehuys, G. R., see Angers, D. A. 444
Miura, T., see Kanetoshi, A. 145
Miyazaki, K.
——, Tanaka, Y. and Saito, M.
Effect of temperature difference between pump and column in molecular weight determination by gel permeation chromatography 357

Moravcová, A., see Matisová, E. 65
Moschidis, M. C.
Silicic acid column chromatography of phosphonolipids. X. Some phosphono analogues of 1-O-alkylethylene glycol 420
Nadeau, P., see Angers, D. A. 444
Nakazawa, H., see Saito, K. 387
Narasimha, R., see Husain, S. 317
Niessen, W. M. A.
——, Bergers, P. J. M., Tjaden, U. R. and Van der Greef, J.
Phase-system switching as an on-line sample pretreatment in the bioanalysis of Mitomycin C using supercritical fluid chromatography 243
Nose, N., see Saito, K. 387
Ogawa, H., see Kanetoshi, A. 145
Ogawa, T., see Terabayashi, T. 432
Ohnishi, K., see Suemitsu, R. 406
Okada, Y., see Koizumi, K. 303
Petsev, N., see Gavrilova, T. B. 73
Phillips, M. W.
——, Subramanian, G. and Cramer, S. M.
Theoretical optimization of operating parameters in non-ideal displacement chromatography 1
Pichot, A., see Liautard, J. 195
Polymeropoulos, C. E., see Kaufman, A. E. 23
Porter, D. W.
——, Banks, M. A., Castranova, V. and Martin, W. G.
Reversed-phase high-performance liquid chromatography technique for taurine quantitation 311
Prasada Raju, N., see Husain, S. 317
Raglione, T. V.
—— and Hartwick, R. A.
On-line high-performance liquid chromatography–post-column reaction–capillary gas chromatography analysis of lipids in biological samples 157
Rüstig, H., see Voelkel, A. 51
Saito, K.
——, Horie, M., Hoshino, Y., Nose, N., Shida, Y., Nakazawa, H. and Fujita, M.
Isolation of virginiamycin-M_1 by droplet counter-current chromatography 387
Saito, M., see Miyazaki, K. 357
Saitoh, E., see Tokuda, H. 109
Sastry, G. S. R., see Husain, S. 317
Schindler, M., see Steuer, W. 253
Sherblom, P. M.
—— and Eganhouse, R. P.
Correlations between octanol–water partition coefficients and reversed-phase high-performance liquid chromatography capacity factors. Chlorobiphenyls and alkylbenzenes 37
Shibamoto, T., see Kallio, H. 392
Shida, Y., see Saito, K. 387
Skarping, G.
——, Willers, S. and Dalene, M.
Determination of cotinine in urine using glass capillary gas chromatography and selective detection, with special reference to the biological monitoring of passive smoking 293
Slominski, B. A.
—— and Campbell, L. D.
Gas chromatographic determination of indoleacetonitriles in rapeseed and *Brassica* vegetables 285
Sri-Widada, J., see Liautard, J. 195
Steuer, W.
——, Schindler, M. and Erni, F.
Gradient elution with normal phases on silica. A comparison between high-performance liquid and supercritical fluid chromatography 253
Stoev, G.
—— and Dentshev, T.
Increasing the reliability of the identification of polymers by pyrolysis gas chromatography 367
Subramanian, G., see Phillips, M. W. 1
Suemitsu, R.
——, Horiuchi, K., Ohnishi, K. and Yanagawase, S.
High-performance liquid chromatographic determination of macrosporin, altersolanol A, alterporriol A, B and C in fermentation of *Alternaria porri* (Ellis) Ciferri 406
Suffet, I. H., see Ibrahim, E. A. 217
Szeleczki, E., see Gazdag, M. 83
Szepesi, G., see Gazdag, M. 83, 95
Szymanowski, J., see Voelkel, A. 51
Takano, S., see Tokuda, H. 109
Tanaka, Y., see Miyazaki, K. 357
Tanimoto, T., see Koizumi, K. 303
Terabayashi, T.
——, Ogawa, T., Kawanishi, Y. and Ishii, J.
In situ mineralization and determination of phosphorus in phospholipids on silica gel sintered thin-layer chromatographic plates 432
Tiebach, R.
—— and Blaas, W.
Direct coupling of a gas chromatograph to an ion trap detector 372
Timossi, A., see Castello, G. 129
Tjaden, U. R., see Niessen, W. M. A. 243
Tokuda, H.
——, Saitoh, E., Kimura, Y. and Takano, S.
Automated analysis of various compounds with a wide range of boiling points by capillary gas chromatography based on retention indices 109

Topalova, I., see Gavrilova, T. B. 73
Van der Greef, J., see Niessen, W. M. A. 243
Vlasenko, E. V., see Gavrilova, T. B. 73
Voelkel, A.
——, Szymanowski, J., Beger, J. and Rüstig, H. Measurement of the polarity of alkyl derivatives of diazapolyoxyethylene ethers by gas chromatography 51
Vučković, G.
——, Malinar, M. J. and Čelap, M. B. Dependence of R_F values of cobalt(III) complexes on the size of diamine chelate ligands on silica gel thin layers 362
Wičar, S.
Sedimentation focusing field-flow fractionation in channels of triangular cross-section 335
Willers, S., see Skarping, G. 293
Wils, E. R. J.
—— and Hulst, A. G. Determination of organophosphorus acids by thermospray liquid chromatography–mass spectrometry 261
Xiao, R., see Zhang, T.-Y. 185
Yanagawase, S., see Suemitsu, R. 406
Zhang, T.-Y.
——, Lee, Y.-W., Fang, Q. C., Xiao, R. and Ito, Y. Preliminary applications of cross-axis synchronous flow-through coil planet centrifuge for large-scale preparative counter-current chromatography 185